

NutriHealers

www.nutrihealers.com

ANTIFUNGUS

CONDICIONES

- Hongos de Uñas

ESPECIES *Copaífera paupera*, *Origanum vulgare*, *Schinus molle*, *Tropaeolum majus*

UBICACIÓN En la costa, sierra y selva del Perú

DESCRIPCION

Copaífera paupera: Árbol de entre 20 y 30 metros de altura, recto de copa globosa. Hojas paripinadas coriáceas alternas, lustrosas, de 3 a 5 cms de largo y 1 a 2 de ancho; inflorescencia terminal racimosa, flores blancas, sésiles, pequeñas, olorosas; fruto legumbre con una a cuatro semillas cubiertas con un arilo anaranjado vivo.

Origanum vulgare: Herbácea de hasta 60cms de alto. Coposa, hojas opuestas, ovales-oviales, pilosas, flores blancas – blanquecinas, aromática.

Schinus molle: Árbol de copa amplia, ramas numerosas y colgantes, flores pequeñas en panículas, frutos en racimo de color rosado. Aroma característico. Contiene aceites esenciales.

Tropaeolum majus: Herbácea, rastrera, trepadora, Tallo carnoso, liso, ramificado, hojas alternas peltado-circulares, flores axilares, tubulares del amarillo al rojo.

Cymbopogon citratus: Herbácea, hojas en macollas compactas, largas, lanceoladas de hasta 1 mt de alto, tallos múltiples cortos, aromática

USOS TRADICIONALES DE LOS INGREDIENTES

Copaífera paupera:

- Micosis: Hongos
- Anti microbiana
- Cáncer
- Cicatrizante
- Herpes
- Psoriasis
- Amigdalitis

Origanum vulgare:

- Reumatismo
- Cólicos menstruales
- Calmante

Schinus molle:

- Antimicrobiano
- Afecciones bronquiales
- Reumatismo



Tropaeolum majus:

- Hongos
- Afecciones cutáneas
- Afecciones respiratorias
- Caspa
- Desinfectante



Cymbopogon citratus:

- Repelente
- Inflamación faríngea
- Anti espasmódico



ESTUDIOS MODERNOS DE LOS INGREDIENTES

Copaifera paupera:

- Antimicótica: Contra hongos
- Anti microbiana
- Cáncer
- Cicatrizante

Origanum vulgare:

- Hongos
- Antioxidante
- Cáncer

Schinus molle:

- Anti microbiano
- Insecticida
- Repelente
- Anti inflamatorio

Tropaeolum majus:

- Anti micótica: Contra los hongos
- Anti microbiana
- Anti tumoral

Cymbopogon citratus:

- Anti Hongos
- Anti microbiano
- Repelente de insectos
- Anti inflamatorio
- Antioxidante

FUENTES

Copaifera paupera:

Planta Med. 2002 Sep;68(9):808-12.

Antimicrobial terpenoids from the oleoresin of the Peruvian medicinal plant *Copaifera paupera*.

Tincusi BM, Jiménez IA, Bazzocchi IL, Moujir LM, Mamani ZA, Barroso JP, Ravelo AG, Hernández BV.

Instituto Universitario de Bio-Orgánica Antonio González, Universidad de La Laguna, La Laguna, Tenerife, Spain.

[ftp://124.42.15.59/ck/2010-](ftp://124.42.15.59/ck/2010-12/165/001/452/991/Antimicrobial%20Terpenoids%20from%20the%20Oleoresin%20of%20the%20Peruvian%20Medicinal%20Plant%20Copaifera%20paupera.pdf)

[12/165/001/452/991/Antimicrobial%20Terpenoids%20from%20the%20Oleoresin%20of%20the%20Peruvian%20Medicinal%20Plant%20Copaifera%20paupera.pdf](ftp://124.42.15.59/ck/2010-12/165/001/452/991/Antimicrobial%20Terpenoids%20from%20the%20Oleoresin%20of%20the%20Peruvian%20Medicinal%20Plant%20Copaifera%20paupera.pdf)

http://www.econbot.org/_about_/06_awards/awards_morton/pdfs/a_herforth.pdf

Origanum vulgare:

Ramírez González, Sandra; López Báez, Orlando; Guzmán Hernández, Tomás; Munguía Ulloa, Sayra; Espinosa Zaragoza, Saúl.

Actividad antifúngica *in vitro* de extractos de *Origanum vulgare* L., *Tradescantia spathacea* Swartz y *Zingiber officinale* Roscoe sobre *Moniliophthora roreri* (Cif & Par) Evans *et ál.*
Tecnología en Marcha, Vol. 24, N.º 2, Abril-Junio 2011, P. 3-17

Brazilian Journal of Microbiology

Print version ISSN 1517-8382

Braz. J. Microbiol. vol.39 no.2 São Paulo Apr./June 2008

<http://dx.doi.org/10.1590/S1517-83822008000200030>

FOOD MICROBIOLOGY

The potential of *Origanum vulgare* L. (Lamiaceae) essential oil in inhibiting the growth of some food-related *Aspergillus* species

Antioxidant and antimicrobial activities of essential oils obtained from oregano (*Origanum vulgare* ssp. *hirtum*) by using different extraction methods.

Karakaya S, El SN, Karagözlü N, Sahin S.

J Med Food. 2011 Jun;14(6):645-52. Epub 2011 Feb 11.

PMID:v21314366

[PubMed - indexed for MEDLINE]

Cancer Lett. 2011 Dec 22;312(2):178-88. doi: 10.1016/j.canlet.2011.08.001. Epub 2011 Aug 26.

β -Caryophyllene oxide inhibits growth and induces apoptosis through the suppression of PI3K/AKT/mTOR/S6K1 pathways and ROS-mediated MAPKs activation.

Park KR, Nam D, Yun HM, Lee SG, Jang HJ, Sethi G, Cho SK, Ahn KS.

College of Oriental Medicine and Institute of Oriental Medicine, Kyung Hee University, Hoegi-Dong Dongdaemun-Gu, Seoul, Republic of Korea.

Nutr Cancer. 2009;61(3):381-9.

Origanum vulgare induces apoptosis in human colon cancer caco2 cells.

Savini J, Arnone R, Catani MV, Avigliano L.

Department of Experimental Medicine and Biochemical Sciences, University of Rome Tor Vergata, Rome 00133, Italy.

J Appl Microbiol. 2007 Oct;103(4):1021-30.

Antifungal effects of essential oils from oregano and fennel on *Sclerotinia sclerotiorum*.

Soylu S, Yigitbas H, Soylu EM, Kurt S.

Mustafa Kemal University, Department of Plant Protection, Faculty of Agriculture, Antakya, Hatay, Turkey.

Schinus molle:

J Ethnopharmacol. 2005 Jan 15;96(3):563-8. Epub 2004 Nov 25.

Screening of antifungal agents using ethanol precipitation and bioautography of medicinal and food plants.

Schmourlo G, Mendonça-Filho RR, Alviano CS, Costa SS.

Núcleo de Pesquisas de Produtos Naturais, Universidade Federal do Rio de Janeiro (UFRJ), Centro de Ciências da Saúde (CCS), Bloco H, Cidade Universitária, Ilha do Fundão, Rio de Janeiro, RJ 21941-590, Brazil.

Cent Afr J Med. 1993 Nov;39(11):231-4.

Antimicrobial activity of essential oil from *Schinus molle* Linn.

Gundidza M.

Department of Pharmacy, Faculty of Medicine, University of Zimbabwe, Harare.

Chemical composition, insecticidal and insect repellent activity of *Schinus molle* L. leaf and fruit essential oils against *Trogoderma granarium* and *Tribolium castaneum*

DOI:10.1080/14786410802346223

Essam Abdel-Sattar^{ab*}, Ahmed A. Zaitoun^c, Mohamed A. Farag^d, Sabah H. El Gayed^b & Fathalla M.H. Harraz^b

Planta Med. 2003 Oct;69(10):893-8.

Isolation of two triterpenoids and a biflavanone with anti-inflammatory activity from *Schinus molle* fruits.

Yueqin Z, Recio MC, Máñez S, Giner RM, Cerdá-Nicolás M, Ríos JL.

Departament de Farmacologia, Facultat de Farmàcia, Universitat de València, Burjassot, Valencia, Spai

Healing activity of ointment the essential oil of the *Schinus molle* L. "molle" in front to the wounds infected in the cattle and in mice

Alba Gonzalez Alex¹, Bonilla Rivera Pablo¹, Arroyo Acevedo Jorge²

¹ Instituto de Ciencias Farmacéuticas y Recursos Naturales "Juan de Dios Guevara" Facultad de Farmacia y Bioquímica. UNMSM. ² Laboratorio de Farmacología. Facultad de Medicina. UNMSM.

Ruiz, J. & Huamani, M. Determinación de la Actividad Antifúngica contra

Candida albicans y *Aspergillus niger* de 10 Plantas Medicinales de 3 Departamentos del Perú

Tropaeolum majus:

[The antimicrobial substance from the Indian cress (*Tropaeolum maius*)].

DANNENBERG H, STICKL H, WENZEL F.

Hoppe Seylers Z Physiol Chem. 1956 Mar 17;303(3-6):248-56. German.

PMID: 13331480 [PubMed - indexed for MEDLINE]

http://www.ucsur.edu.pe/documentos/cientifica_06.pdf

Planta Med. 1995 Jun;61(3):233-6.

In vitro and in vivo antitumor activity of benzyl isothiocyanate: a natural product from *Tropaeolum majus*.

Pintão AM, Pais MS, Coley H, Kelland LR, Judson IR.

Departamento de Biologia Vegetal, Faculdade de Ciências de Lisboa, Portugal.

Cymbopogon citratus

In vitro antifungal, anti-elastase and anti-keratinase activity of essential oils of *Cinnamomum*-, *Syzygium*- and *Cymbopogon*-species against *Aspergillus fumigatus* and *Trichophyton rubrum*.

Khan MS, Ahmad I.

Phytomedicine. 2011 Dec 15;19(1):48-55. Epub 2011 Sep 3.

PMID: 21893402

Effectiveness of *Cymbopogon citratus* L. essential oil to inhibit the growth of some filamentous fungi and yeasts.

Irkin R, Korukluoglu M.

J Med Food. 2009 Feb;12(1):193-7.

PMID: 19298215 [PubMed - indexed for MEDLINE]

Chemical composition and antimicrobial activity of *Cymbopogon citratus* and *Cymbopogon giganteus* essential oils alone and in combination.

Bassolé IH, Lamien-Meda A, Bayala B, Obame LC, Ilboudo AJ, Franz C, Novak J, Nebié RC, Dicko MH.

Phytomedicine. 2011 Sep 15;18(12):1070-4. Epub 2011 Jun 12.

PMID: 21665450 [PubMed - indexed for MEDLINE]

Repellent, larvicidal and pupicidal properties of essential oils and their formulations against the housefly, *Musca domestica*.

Kumar P, Mishra S, Malik A, Satya S.

Med Vet Entomol. 2011 Sep;25(3):302-10. doi: 10.1111/j.1365-2915.2011.00945.x. Epub 2011 Feb 21.

PMID: 21338379 [PubMed - indexed for MEDLINE]

Suppression of oxidative stress and pro-inflammatory mediators by *Cymbopogon citratus* D. Stapf extract in lipopolysaccharide stimulated murine alveolar macrophages.

Tiwari M, Dwivedi UN, Kakkar P.

Food Chem Toxicol. 2010 Oct;48(10):2913-9. Epub 2010 Jul 22.

PMID: 20655974 [PubMed - indexed for MEDLINE]

Diccionario enciclopédico de Plantas Útiles del Perú Antonio Brack Egg, PNUD – Centro de Estudios Regionales Andinos Bartolomé de las Casas

All information is for informational purposes only, and not intended to diagnose, treat, cure, mitigate, or prevent any disease. No statements have been evaluated by the FDA. We always suggest talking to your physician concerning any questions you may have about supplement/drug interactions.

Extended Disclaimer [Click Here](#)

NutritHealers